

**GENERATION OF PLANTS WITH ALTERED OIL CONTENT****REFERENCE TO RELATED APPLICATIONS**

**[0001]** This application claims priority to U.S. provisional application 60/530,799 filed December 17, 2003, the contents of which are hereby incorporated by reference.

**BACKGROUND OF THE INVENTION**

**[0002]** The ability to manipulate the composition of crop seeds, particularly the content and composition of seed oils, has important applications in the agricultural industries, relating both to processed food oils and to oils for animal feeding. Seeds of agricultural crops contain a variety of valuable constituents, including oil, protein and starch. Industrial processing can separate some or all of these constituents for individual sale in specific applications. For instance, nearly 60% of the U.S. soybean crop is crushed by the soy processing industry. Soy processing yields purified oil, which is sold at high value, while the remainder is sold principally for lower value livestock feed (U.S. Soybean Board, 2001 Soy Stats). Canola seed is crushed to produce oil and the co-product canola meal (Canola Council of Canada). Nearly 20% of the 1999/2000 U.S. corn crop was industrially refined, primarily for production of starch, ethanol and oil (Corn Refiners Association). Thus, it is often desirable to maximize oil content of seeds. For instance, for processed oilseeds such as soy and canola, increasing the absolute oil content of the seed will increase the value of such grains. For processed corn it may be desired to either increase or decrease oil content, depending on utilization of other major constituents. Decreasing oil may improve the quality of isolated starch by reducing undesired flavors associated with oil oxidation. Alternatively, in ethanol production, where flavor is unimportant, increasing oil content may increase overall value. In many fed grains, such as corn and wheat, it is desirable to increase seed oil content, because oil has higher energy content than other seed constituents such as carbohydrate. Oilseed processing, like most grain processing businesses, is a capital-intensive business; thus small shifts in the distribution of products from the low valued components to the high value oil component can have substantial economic impacts for grain processors.

**[0003]** Biotechnological manipulation of oils can provide compositional alteration and improvement of oil yield. Compositional alterations include high oleic soybean and corn oil (U.S. Pat Nos 6,229,033 and 6,248,939), and laurate-containing seeds (U.S. Pat No 5,639,790), among others. Work in compositional alteration has predominantly focused on processed oilseeds but has been readily extendable to non-oilseed crops, including corn. While there is considerable interest in increasing oil content, the only currently practiced biotechnology in this area is High-Oil Corn (HOC) technology (DuPont, U.S. PAT NO: 5,704,160). HOC employs high oil pollinators developed by classical selection

breeding along with elite (male-sterile) hybrid females in a production system referred to as TopCross. The TopCross High Oil system raises harvested grain oil content in maize from about 3.5% to about 7%, improving the energy content of the grain.

[0004] While it has been fruitful, the HOC production system has inherent limitations. First, the system of having a low percentage of pollinators responsible for an entire field's seed set contains inherent risks, particularly in drought years. Second, oil contents in current HOC fields have plateaued at about 9% oil. Finally, high-oil corn is not primarily a biochemical change, but rather an anatomical mutant (increased embryo size) that has the indirect result of increasing oil content. For these reasons, an alternative high oil strategy, particularly one that derives from an altered biochemical output, would be especially valuable.

[0005] The most obvious target crops for the processed oil market are soy and rapeseed, and a large body of commercial work (e.g., U.S. Pat No: 5,952,544; PCT application WO9411516) demonstrates that *Arabidopsis* is an excellent model for oil metabolism in these crops. Biochemical screens of seed oil composition have identified *Arabidopsis* genes for many critical biosynthetic enzymes and have led to identification of agronomically important gene orthologs. For instance, screens using chemically mutagenized populations have identified lipid mutants whose seeds display altered fatty acid composition (Lemieux et al., 1990; James and Dooner, 1990). T-DNA mutagenesis screens (Feldmann et al., 1989) that detected altered fatty acid composition identified the omega 3 desaturase (*FAD3*) and delta-12 desaturase (*FAD2*) genes (U.S. Pat No 5952544; Yadav et al., 1993; Okuley et al., 1994). A screen which focused on oil content rather than oil quality, analyzed chemically-induced mutants for wrinkled seeds or altered seed density, from which altered seed oil content was inferred (Focks and Benning, 1998). Another screen, designed to identify enzymes involved in production of very long chain fatty acids, identified a mutation in the gene encoding a diacylglycerol acyltransferase (DGAT) as being responsible for reduced triacyl glycerol accumulation in seeds (Katavic V et al, 1995). It was further shown that seed-specific over-expression of the DGAT cDNA was associated with increased seed oil content (Jako et al., 2001).

[0006] Activation tagging in plants refers to a method of generating random mutations by insertion of a heterologous nucleic acid construct comprising regulatory sequences (e.g., an enhancer) into a plant genome. The regulatory sequences can act to enhance transcription of one or more native plant genes; accordingly, activation tagging is a fruitful method for generating gain-of-function, generally dominant mutants (see, e.g., Hayashi et al., 1992; Weigel D et al. 2000). The inserted construct provides a molecular tag for rapid identification of the native plant whose mis-expression causes the mutant phenotype.

Activation tagging may also cause loss-of-function phenotypes. The insertion may result in disruption of a native plant gene, in which case the phenotype is generally recessive.

[0007] Activation tagging has been used in various species, including tobacco and *Arabidopsis*, to identify many different kinds of mutant phenotypes and the genes associated with these phenotypes (Wilson *et al.*, 1996, Schaffer *et al.*, 1998, Fridborg *et al.*, 1999; Kardailsky *et al.*, 1999; Christensen S *et al.* 1998).

### SUMMARY OF THE INVENTION

[0008] The invention provides a transgenic plant having a high oil phenotype. The transgenic plant comprises a transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO32.3 polypeptide comprising the amino acid sequence of SEQ ID NO:2, or an ortholog thereof. In preferred embodiments, the transgenic plant is selected from the group consisting of rapeseed, soy, corn, sunflower, cotton, cocoa, safflower, oil palm, coconut palm, flax, castor and peanut. The invention also provides a method of producing oil comprising growing the transgenic plant and recovering oil from said plant. The invention further provides a method of generating a plant having a high oil phenotype by identifying a plant that has an allele in its HIO30.4 gene that results in increased oil content compared to plants lacking the allele and generating progeny of the identified plant, wherein the generated progeny inherit the allele and have the high oil phenotype.

[0009] The transgenic plant of the invention is produced by a method that comprises introducing into progenitor cells of the plant a plant transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO32.3 polypeptide, and growing the transformed progenitor cells to produce a transgenic plant, wherein the HIO32.3 polynucleotide sequence is expressed causing the high oil phenotype.

### DETAILED DESCRIPTION OF THE INVENTION

#### Definitions

[0010] Unless otherwise indicated, all technical and scientific terms used herein have the same meaning as they would to one skilled in the art of the present invention.

Practitioners are particularly directed to Sambrook *et al.*, 1989, and Ausubel FM *et al.*, 1993, for definitions and terms of the art. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary.

[0011] As used herein, the term "vector" refers to a nucleic acid construct designed for transfer between different host cells. An "expression vector" refers to a vector that has the

ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art.

[0012] A "heterologous" nucleic acid construct or sequence has a portion of the sequence that is not native to the plant cell in which it is expressed. Heterologous, with respect to a control sequence refers to a control sequence (*i.e.* promoter or enhancer) that does not function in nature to regulate the same gene the expression of which it is currently regulating. Generally, heterologous nucleic acid sequences are not endogenous to the cell or part of the genome in which they are present, and have been added to the cell, by infection, transfection, microinjection, electroporation, or the like. A "heterologous" nucleic acid construct may contain a control sequence/DNA coding sequence combination that is the same as, or different from a control sequence/DNA coding sequence combination found in the native plant.

[0013] As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, which may or may not include regions preceding and following the coding region, *e.g.* 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons) and non-transcribed regulatory sequence.

[0014] As used herein, "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

[0015] As used herein, the term "gene expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation; accordingly, "expression" may refer to either a polynucleotide or polypeptide sequence, or both. Sometimes, expression of a polynucleotide sequence will not lead to protein translation. "Over-expression" refers to increased expression of a polynucleotide and/or polypeptide sequence relative to its expression in a wild-type (or other reference [*e.g.*, non-transgenic]) plant and may relate to a naturally-occurring or non-naturally occurring sequence. "Ectopic expression" refers to expression at a time, place, and/or increased level that does not naturally occur in the non-altered or wild-type plant. "Under-expression" refers to decreased expression of a polynucleotide and/or polypeptide sequence, generally of an endogenous gene, relative to

its expression in a wild-type plant. The terms "mis-expression" and "altered expression" encompass over-expression, under-expression, and ectopic expression.

[0016] The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

[0017] As used herein, a "plant cell" refers to any cell derived from a plant, including cells from undifferentiated tissue (*e.g.*, callus) as well as plant seeds, pollen, progagules and embryos.

[0018] As used herein, the terms "native" and "wild-type" relative to a given plant trait or phenotype refers to the form in which that trait or phenotype is found in the same variety of plant in nature.

[0019] As used herein, the term "modified" regarding a plant trait, refers to a change in the phenotype of a transgenic plant relative to the similar non-transgenic plant. An "interesting phenotype (trait)" with reference to a transgenic plant refers to an observable or measurable phenotype demonstrated by a T1 and/or subsequent generation plant, which is not displayed by the corresponding non-transgenic (*i.e.*, a genotypically similar plant that has been raised or assayed under similar conditions). An interesting phenotype may represent an improvement in the plant or may provide a means to produce improvements in other plants. An "improvement" is a feature that may enhance the utility of a plant species or variety by providing the plant with a unique and/or novel quality. An "altered oil content phenotype" refers to measurable phenotype of a genetically modified plant, where the plant displays a statistically significant increase or decrease in overall oil content (*i.e.*, the percentage of seed mass that is oil), as compared to the similar, but non-modified plant. A high oil phenotype refers to an increase in overall oil content.

[0020] As used herein, a "mutant" polynucleotide sequence or gene differs from the corresponding wild type polynucleotide sequence or gene either in terms of sequence or expression, where the difference contributes to a modified plant phenotype or trait. Relative to a plant or plant line, the term "mutant" refers to a plant or plant line which has a modified plant phenotype or trait, where the modified phenotype or trait is associated with the modified expression of a wild type polynucleotide sequence or gene.

[0021] As used herein, the term "T1" refers to the generation of plants from the seed of T0 plants. The T1 generation is the first set of transformed plants that can be selected by application of a selection agent, *e.g.*, an antibiotic or herbicide, for which the transgenic plant contains the corresponding resistance gene. The term "T2" refers to the generation of plants by self-fertilization of the flowers of T1 plants, previously selected as being transgenic. T3 plants are generated from T2 plants, etc. As used herein, the "direct progeny" of a given plant derives from the seed (or, sometimes, other tissue) of that plant and is in the immediately subsequent generation; for instance, for a given lineage, a T2 plant is the direct progeny of a T1 plant. The "indirect progeny" of a given plant derives from the seed (or other tissue) of the direct progeny of that plant, or from the seed (or other tissue) of subsequent generations in that lineage; for instance, a T3 plant is the indirect progeny of a T1 plant.

[0022] As used herein, the term "plant part" includes any plant organ or tissue, including, without limitation, seeds, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. Plant cells can be obtained from any plant organ or tissue and cultures prepared therefrom. The class of plants which can be used in the methods of the present invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledenous and dicotyledenous plants.

[0023] As used herein, "transgenic plant" includes a plant that comprises within its genome a heterologous polynucleotide. The heterologous polynucleotide can be either stably integrated into the genome, or can be extra-chromosomal. Preferably, the polynucleotide of the present invention is stably integrated into the genome such that the polynucleotide is passed on to successive generations. A plant cell, tissue, organ, or plant into which the heterologous polynucleotides have been introduced is considered "transformed", "transfected", or "transgenic". Direct and indirect progeny of transformed plants or plant cells that also contain the heterologous polynucleotide are also considered transgenic.

#### **Identification of Plants with an Altered Oil Content Phenotype**

[0024] We used an *Arabidopsis* activation tagging (ACTTAG) screen to identify the association between the gene we have designated "HIO32.3," (At3g47720; GI#18408490 encoding protein T23J7.50 (GI#15228235), and an altered oil content phenotype (specifically, a high oil phenotype). Briefly, and as further described in the Examples, a large number of *Arabidopsis* plants were mutated by transformation with the pSKI015 vector, which comprises a T-DNA from the Ti plasmid of *Agrobacterium tumefaciens*, a

viral enhancer element, and a selectable marker gene (Weigel *et al*, 2000). When the T-DNA inserts into the genome of transformed plants, the enhancer element can cause up-regulation genes in the vicinity, generally within about 10 kilobase (kb) of the enhancers. To identify transgenic plants, T1 plants were exposed to the selective agent in order to specifically recover transformed plants that expressed the selectable marker and harboring the T-DNA. T2 seed was harvested from these plants. Lipids were extracted from of about 15-20 T2 seeds. Gas chromatography (GC) analysis was performed to determine fatty acid content and composition of seed samples.

[0025] An *Arabidopsis* line that showed a high-oil phenotype was identified, wherein oil content (i.e., fatty acids) constituted about 36.3% of seed mass compared to an average oil content of about 27.0% for seed from other plants grown and analyzed at the same time (a 34 % in oil). The association of the HIO32.3 gene with the high oil phenotype was discovered by identifying the site of T-DNA insertion and, as shown in the Examples, demonstrating genetic co-segregation of the high oil phenotype and the presence of the T-DNA. Accordingly, HIO32.3 genes and/or polypeptides may be employed in the development of genetically modified plants having a modified oil content phenotype ("a HIO32.3 phenotype"). HIO32.3 genes may be used in the generation of oilseed crops that provide improved oil yield from oilseed processing and in the generation of feed grain crops that provide increased energy for animal feeding. HIO32.3 genes may further be used to increase the oil content of specialty oil crops, in order to augment yield of desired unusual fatty acids. Transgenic plants that have been genetically modified to express HIO32.3 can be used in the production of oil, wherein the transgenic plants are grown, and oil is obtained from plant parts (e.g. seed) using standard methods.

#### HIO32.3 Nucleic Acids and Polypeptides

[0026] *Arabidopsis* HIO32.3 nucleic acid (genomic DNA) sequence is provided in SEQ ID NO:1 and in Genbank entry GI#18408490. The corresponding protein sequence is provided in SEQ ID NO:2 and in GI#15228235. Nucleic acids and/or proteins that are orthologs or paralogs of *Arabidopsis* HIO32.3, are described in Example 3 below.

[0027] As used herein, the term "HIO32.3 polypeptide" refers to a full-length HIO32.3 protein or a fragment, derivative (variant), or ortholog thereof that is "functionally active," meaning that the protein fragment, derivative, or ortholog exhibits one or more of the functional activities associated with the polypeptide of SEQ ID NO:2. In one preferred embodiment, a functionally active HIO32.3 polypeptide causes an altered oil content phenotype when mis-expressed in a plant. In a further preferred embodiment, mis-expression of the HIO32.3 polypeptide causes a high oil phenotype in a plant. In another embodiment, a functionally active HIO32.3 polypeptide is capable of rescuing defective

(including deficient) endogenous HIO32.3 activity when expressed in a plant or in plant cells; the rescuing polypeptide may be from the same or from a different species as that with defective activity. In another embodiment, a functionally active fragment of a full length HIO32.3 polypeptide (i.e., a native polypeptide having the sequence of SEQ ID NO:2 or a naturally occurring ortholog thereof) retains one or more of the biological properties associated with the full-length HIO32.3 polypeptide, such as signaling activity, binding activity, catalytic activity, or cellular or extra-cellular localizing activity. A HIO32.3 fragment preferably comprises a HIO32.3 domain, such as a C- or N-terminal or catalytic domain, among others, and preferably comprises at least 10, preferably at least 20, more preferably at least 25, and most preferably at least 50 contiguous amino acids of a HIO32.3 protein. Functional domains can be identified using the PFAM program (Bateman A et al., 1999 Nucleic Acids Res 27:260-262). Functionally active variants of full-length HIO32.3 polypeptides or fragments thereof include polypeptides with amino acid insertions, deletions, or substitutions that retain one or more of the biological properties associated with the full-length HIO32.3 polypeptide. In some cases, variants are generated that change the post-translational processing of a HIO32.3 polypeptide. For instance, variants may have altered protein transport or protein localization characteristics or altered protein half-life compared to the native polypeptide.

[0028] As used herein, the term "HIO32.3 nucleic acid" encompasses nucleic acids with the sequence provided in or complementary to the sequence provided in SEQ ID NO:1, as well as functionally active fragments, derivatives, or orthologs thereof. A HIO32.3 nucleic acid of this invention may be DNA, derived from genomic DNA or cDNA, or RNA.

[0029] In one embodiment, a functionally active HIO32.3 nucleic acid encodes or is complementary to a nucleic acid that encodes a functionally active HIO32.3 polypeptide. Included within this definition is genomic DNA that serves as a template for a primary RNA transcript (i.e., an mRNA precursor) that requires processing, such as splicing, before encoding the functionally active HIO32.3 polypeptide. A HIO32.3 nucleic acid can include other non-coding sequences, which may or may not be transcribed; such sequences include 5' and 3' UTRs, polyadenylation signals and regulatory sequences that control gene expression, among others, as are known in the art. Some polypeptides require processing events, such as proteolytic cleavage, covalent modification, etc., in order to become fully active. Accordingly, functionally active nucleic acids may encode the mature or the pre-processed HIO32.3 polypeptide, or an intermediate form. A HIO32.3 polynucleotide can also include heterologous coding sequences, for example, sequences that encode a marker included to facilitate the purification of the fused polypeptide, or a transformation marker.



[0030] In another embodiment, a functionally active HIO32.3 nucleic acid is capable of being used in the generation of loss-of-function HIO32.3 phenotypes, for instance, via antisense suppression, co-suppression, etc.

[0031] In one preferred embodiment, a HIO32.3 nucleic acid used in the methods of this invention comprises a nucleic acid sequence that encodes or is complementary to a sequence that encodes a HIO32.3 polypeptide having at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95% or more sequence identity to the polypeptide sequence presented in SEQ ID NO:2.

[0032] In another embodiment a HIO32.3 polypeptide of the invention comprises a polypeptide sequence with at least 50% or 60% identity to the HIO32.3 polypeptide sequence of SEQ ID NO:2, and may have at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the HIO32.3 polypeptide sequence of SEQ ID NO:2. In another embodiment, a HIO32.3 polypeptide comprises a polypeptide sequence with at least 50%, 60%, 70%, 80%, 85%, 90% or 95% or more sequence identity to a functionally active fragment of the polypeptide presented in SEQ ID NO:2. In yet another embodiment, a HIO32.3 polypeptide comprises a polypeptide sequence with at least 50%, 60%, 70%, 80%, or 90% identity to the polypeptide sequence of SEQ ID NO:2 over its entire length.

[0033] In another aspect, a HIO32.3 polynucleotide sequence is at least 50% to 60% identical over its entire length to the HIO32.3 nucleic acid sequence presented as SEQ ID NO:1, or nucleic acid sequences that are complementary to such a HIO32.3 sequence, and may comprise at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the HIO32.3 sequence presented as SEQ ID NO:1 or a functionally active fragment thereof, or complementary sequences.

[0034] As used herein, "percent (%) sequence identity" with respect to a specified subject sequence, or a specified portion thereof, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1990) 215:403-410) with search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A "% identity value" is determined by the number of matching identical nucleotides or amino acids divided by the

sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation. A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

[0035] Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that selectively hybridize to the nucleic acid sequence of SEQ ID NO:1. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are well known (see, *e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of SEQ ID NO:1 under stringent hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1 h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate). In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS. Alternatively, low stringency conditions can be used that comprise: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon

sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

[0036] As a result of the degeneracy of the genetic code, a number of polynucleotide sequences encoding a HIO32.3 polypeptide can be produced. For example, codons may be selected to increase the rate at which expression of the polypeptide occurs in a particular host species, in accordance with the optimum codon usage dictated by the particular host organism (see, e.g., Nakamura et al, 1999). Such sequence variants may be used in the methods of this invention.

[0037] The methods of the invention may use orthologs of the *Arabidopsis* HIO32.3. Methods of identifying the orthologs in other plant species are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Arabidopsis*, may correspond to multiple genes (paralogs) in another. As used herein, the term "orthologs" encompasses paralogs. When sequence data is available for a particular plant species, orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. Nucleic acid hybridization methods may also be used to find orthologous genes and are preferred when sequence data are not available. Degenerate PCR and screening of cDNA or genomic DNA libraries are common methods for finding related gene sequences and are well known in the art (see, e.g., Sambrook, 1989; Dieffenbach and Dveksler, 1995). For instance, methods for generating a cDNA library from the plant species of interest and probing the library with partially homologous gene probes are described in Sambrook *et al.* A highly conserved portion of the *Arabidopsis* HIO32.3 coding sequence may be used as a probe. HIO32.3 ortholog nucleic acids may hybridize to the nucleic acid of SEQ ID NO:1 under high, moderate, or low stringency conditions. After amplification or isolation of a segment of a

putative ortholog, that segment may be cloned and sequenced by standard techniques and utilized as a probe to isolate a complete cDNA or genomic clone. Alternatively, it is possible to initiate an EST project to generate a database of sequence information for the plant species of interest. In another approach, antibodies that specifically bind known HIO32.3 polypeptides are used for ortholog isolation (see, e.g., Harlow and Lane, 1988, 1999). Western blot analysis can determine that a HIO32.3 ortholog (i.e., an orthologous protein) is present in a crude extract of a particular plant species. When reactivity is observed, the sequence encoding the candidate ortholog may be isolated by screening expression libraries representing the particular plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in Sambrook, *et al.*, 1989. Once the candidate ortholog(s) are identified by any of these means, candidate orthologous sequence are used as bait (the "query") for the reverse BLAST against sequences from *Arabidopsis* or other species in which HIO32.3 nucleic acid and/or polypeptide sequences have been identified.

[0038] HIO32.3 nucleic acids and polypeptides may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR), as previously described, are well known in the art. Alternatively, nucleic acid sequence may be synthesized. Any known method, such as site directed mutagenesis (Kunkel TA *et al.*, 1991), may be used to introduce desired changes into a cloned nucleic acid.

[0039] In general, the methods of the invention involve incorporating the desired form of the HIO32.3 nucleic acid into a plant expression vector for transformation of in plant cells, and the HIO32.3 polypeptide is expressed in the host plant.

[0040] An isolated HIO32.3 nucleic acid molecule is other than in the form or setting in which it is found in nature and is identified and separated from least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the HIO32.3 nucleic acid. However, an isolated HIO32.3 nucleic acid molecule includes HIO32.3 nucleic acid molecules contained in cells that ordinarily express HIO32.3 where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

### **Generation of Genetically Modified Plants with an Altered Oil Content**

#### **Phenotype**

[0041] HIO32.3 nucleic acids and polypeptides may be used in the generation of genetically modified plants having a modified oil content phenotype. As used herein, a "modified oil content phenotype" may refer to modified oil content in any part of the plant; the modified oil content is often observed in seeds. In a preferred embodiment,

altered expression of the HIO32.3 gene in a plant is used to generate plants with a high oil phenotype.

[0042] The methods described herein are generally applicable to all plants. Although activation tagging and gene identification is carried out in *Arabidopsis*, the HIO32.3 gene (or an ortholog, variant or fragment thereof) may be expressed in any type of plant. In a preferred embodiment, the invention is directed to oil-producing plants, which produce and store triacylglycerol in specific organs, primarily in seeds. Such species include soybean (*Glycine max*), rapeseed and canola (including *Brassica napus*, *B. campestris*), sunflower (*Helianthus annuus*), cotton (*Gossypium hirsutum*), corn (*Zea mays*), cocoa (*Theobroma cacao*), safflower (*Carthamus tinctorius*), oil palm (*Elaeis guineensis*), coconut palm (*Cocos nucifera*), flax (*Linum usitatissimum*), castor (*Ricinus communis*) and peanut (*Arachis hypogaea*). The invention may also be directed to fruit- and vegetable-bearing plants, grain-producing plants, nut-producing plants, rapid cycling *Brassica* species, alfalfa (*Medicago sativa*), tobacco (*Nicotiana*), turfgrass (*Poaceae* family), other forage crops, and wild species that may be a source of unique fatty acids.

[0043] The skilled artisan will recognize that a wide variety of transformation techniques exist in the art, and new techniques are continually becoming available. Any technique that is suitable for the target host plant can be employed within the scope of the present invention. For example, the constructs can be introduced in a variety of forms including, but not limited to as a strand of DNA, in a plasmid, or in an artificial chromosome. The introduction of the constructs into the target plant cells can be accomplished by a variety of techniques, including, but not limited to *Agrobacterium*-mediated transformation, electroporation, microinjection, microprojectile bombardment calcium-phosphate-DNA co-precipitation or liposome-mediated transformation of a heterologous nucleic acid. The transformation of the plant is preferably permanent, *i.e.* by integration of the introduced expression constructs into the host plant genome, so that the introduced constructs are passed onto successive plant generations. Depending upon the intended use, a heterologous nucleic acid construct comprising an HIO32.3 polynucleotide may encode the entire protein or a biologically active portion thereof.

[0044] In one embodiment, binary Ti-based vector systems may be used to transfer polynucleotides. Standard *Agrobacterium* binary vectors are known to those of skill in the art, and many are commercially available (e.g., pBI121 Clontech Laboratories, Palo Alto, CA).

[0045] The optimal procedure for transformation of plants with *Agrobacterium* vectors will vary with the type of plant being transformed. Exemplary methods for *Agrobacterium*-mediated transformation include transformation of explants of hypocotyl, shoot tip, stem or leaf tissue, derived from sterile seedlings and/or plantlets. Such transformed plants may be reproduced sexually, or by cell or tissue culture. *Agrobacterium* transformation has been previously described for a large number of different types of plants and methods for such transformation may be found in the scientific literature. Of particular relevance are methods to transform commercially important crops, such as rapeseed (De Block et al., 1989), sunflower (Everett et al., 1987), and soybean (Christou et al., 1989; Kline et al., 1987).

[0046] Expression (including transcription and translation) of HIO32.3 may be regulated with respect to the level of expression, the tissue type(s) where expression takes place and/or developmental stage of expression. A number of heterologous regulatory sequences (e.g., promoters and enhancers) are available for controlling the expression of a HIO32.3 nucleic acid. These include constitutive, inducible and regulatable promoters, as well as promoters and enhancers that control expression in a tissue- or temporal-specific manner. Exemplary constitutive promoters include the raspberry E4 promoter (U.S. Patent Nos. 5,783,393 and 5,783,394), the 35S CaMV (Jones JD *et al.*, 1992), the CsVMV promoter (Verd  guer B *et al.*, 1998) and the melon actin promoter (published PCT application WO0056863). Exemplary tissue-specific promoters include the tomato E4 and E8 promoters (U.S. Patent No. 5,859,330) and the tomato 2AII gene promoter (Van Haaren MJJ *et al.*, 1993).

[0047] In one preferred embodiment, HIO32.3 expression is under control of regulatory sequences from genes whose expression is associated with early seed and/or embryo development. Legume genes whose promoters are associated with early seed and embryo development include *V. faba legumin* (Baumlein et al., 1991, Mol Gen Genet 225:121-8; Baumlein et al., 1992, Plant J 2:233-9), *V. faba usp* (Fiedler et al., 1993, Plant Mol Biol 22:669-79), pea *convicilin* (Bown et al., 1988, Biochem J 251:717-26), pea *lectin* (dePater et al., 1993, Plant Cell 5:877-86), *P. vulgaris beta phaseolin* (Bustos et al., 1991, EMBO J 10:1469-79), *P. vulgaris DLEC2* and *PHS* [beta] (Bobb et al., 1997, Nucleic Acids Res 25:641-7), and soybean *beta-Conglycinin*, 7S storage protein (Chamberland et al., 1992, Plant Mol Biol 19:937-49). Cereal genes whose promoters are associated with early seed and embryo development include rice *glutelin* ("GluA-3," Yoshihara and Takaiwa, 1996, Plant Cell Physiol 37:107-11; "GluB-1," Takaiwa et al., 1996, Plant Mol Biol 30:1207-21; Washida et al., 1999, Plant Mol Biol 40:1-12; "Gt3," Leisy et al., 1990, Plant Mol Biol 14:41-50), rice *prolamin* (Zhou & Fan, 1993, Transgenic Res 2:141-6), wheat *prolamin* (Hammond-Kosack et al., 1993, EMBO J 12:545-54), maize *zein* (Z4, Matzke et al., 1990,

Plant Mol Biol 14:323-32), and barley *B-hordeins* (Entwistle et al., 1991, Plant Mol Biol 17:1217-31). Other genes whose promoters are associated with early seed and embryo development include oil palm GLO7A (7S globulin, Morcillo et al., 2001, Physiol Plant 112:233-243), *Brassica napus napin*, 2S storage protein, and napA gene (Josefsson et al., 1987, J Biol Chem 262:12196-201; Stalberg et al., 1993, Plant Mol Biol 1993 23:671-83; Ellerstrom et al., 1996, Plant Mol Biol 32:1019-27), *Brassica napus oleosin* (Keddie et al., 1994, Plant Mol Biol 24:327-40), *Arabidopsis oleosin* (Plant et al., 1994, Plant Mol Biol 25:193-205), *Arabidopsis* FAE1 (Rossak et al., 2001, Plant Mol Biol 46:717-25), *Canavalia gladiata* conA (Yamamoto et al., 1995, Plant Mol Biol 27:729-41), and *Catharanthus roseus* strictosidine synthase (Str, Ouwerkerk and Memelink, 1999, Mol Gen Genet 261:635-43). In another preferred embodiment, regulatory sequences from genes expressed during oil biosynthesis are used (see, e.g., US Pat No: 5,952, 544). Alternative promoters are from plant storage protein genes (Bevan et al., 1993, Philos Trans R Soc Lond B Biol Sci 342:209-15).

[0048] In yet another aspect, in some cases it may be desirable to inhibit the expression of endogenous HIO32.3 in a host cell. Exemplary methods for practicing this aspect of the invention include, but are not limited to antisense suppression (Smith, *et al.*, 1988; van der Krol et al., 1988); co-suppression (Napoli, *et al.*, 1990); ribozymes (PCT Publication WO 97/10328); and combinations of sense and antisense (Waterhouse, *et al.*, 1998). Methods for the suppression of endogenous sequences in a host cell typically employ the transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well as non-coding regions of the endogenous sequence. Antisense inhibition may use the entire cDNA sequence (Sheehy et al., 1988), a partial cDNA sequence including fragments of 5' coding sequence, (Cannon et al., 1990), or 3' non-coding sequences (Ch'ng et al., 1989). Cosuppression techniques may use the entire cDNA sequence (Napoli et al., 1990; van der Krol et al., 1990), or a partial cDNA sequence (Smith et al., (1990).

[0049] Standard molecular and genetic tests may be performed to further analyze the association between a gene and an observed phenotype. Exemplary techniques are described below.

#### DNA/RNA analysis

[0050] The stage- and tissue-specific gene expression patterns in mutant versus wild-type lines may be determined, for instance, by in situ hybridization. Analysis of the methylation status of the gene, especially flanking regulatory regions, may be performed. Other suitable techniques include overexpression, ectopic expression, expression in other plant

species and gene knock-out (reverse genetics, targeted knock-out, viral induced gene silencing [VIGS, see Baulcombe D, 1999]).

[0051] In a preferred application expression profiling, generally by microarray analysis, is used to simultaneously measure differences or induced changes in the expression of many different genes. Techniques for microarray analysis are well known in the art (Schena M *et al.*, Science (1995) 270:467-470; Baldwin D *et al.*, 1999; Dangond F, Physiol Genomics (2000) 2:53-58; van Hal NL *et al.*, J Biotechnol (2000) 78:271-280; Richmond T and Somerville S, Curr Opin Plant Biol (2000) 3:108-116). Expression profiling of individual tagged lines may be performed. Such analysis can identify other genes that are coordinately regulated as a consequence of the overexpression of the gene of interest, which may help to place an unknown gene in a particular pathway.

#### **Gene Product Analysis**

[0052] Analysis of gene products may include recombinant protein expression, antisera production, immunolocalization, biochemical assays for catalytic or other activity, analysis of phosphorylation status, and analysis of interaction with other proteins via yeast two-hybrid assays.

#### **Pathway Analysis**

[0053] Pathway analysis may include placing a gene or gene product within a particular biochemical, metabolic or signaling pathway based on its mis-expression phenotype or by sequence homology with related genes. Alternatively, analysis may comprise genetic crosses with wild-type lines and other mutant lines (creating double mutants) to order the gene in a pathway, or determining the effect of a mutation on expression of downstream "reporter" genes in a pathway.

#### **Generation of Mutated Plants with an Altered Oil Content Phenotype**

[0054] The invention further provides a method of identifying non-transgenic plants that have mutations in or an allele of endogenous HIO32.3 that confer a HIO32.3 phenotype to these plants and their progeny. In one method, called "TILLING" (for targeting induced local lesions in genomes), mutations are induced in the seed of a plant of interest, for example, using EMS treatment. The resulting plants are grown and self-fertilized, and the progeny are used to prepare DNA samples. HIO32.3 -specific PCR is used to identify whether a mutated plant has a HIO32.3 mutation. Plants having HIO32.3 mutations may then be tested for altered oil content, or alternatively, plants may be tested for altered oil content, and then HIO32.3 -specific PCR is used to determine whether a plant having



altered oil content has a mutated HIO32.3 gene. TILLING can identify mutations that may alter the expression of specific genes or the activity of proteins encoded by these genes (see Colbert et al (2001) Plant Physiol 126:480-484; McCallum et al (2000) Nature Biotechnology 18:455-457).

[0055] In another method, a candidate gene/Quantitative Trait Locus (QTLs) approach can be used in a marker-assisted breeding program to identify alleles of or mutations in the HIO32.3 gene or orthologs of HIO32.3 that may confer altered oil content (see Bert et al., Theor Appl Genet. 2003 Jun;107(1):181-9; and Lionneton et al, Genome. 2002 Dec;45(6):1203-15). Thus, in a further aspect of the invention, a HIO32.3 nucleic acid is used to identify whether a plant having altered oil content has a mutation in endogenous HIO32.3 or has a particular allele that causes altered oil content.

[0056] While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention. All publications cited herein are expressly incorporated herein by reference for the purpose of describing and disclosing compositions and methodologies that might be used in connection with the invention. All cited patents, patent applications, and sequence information in referenced public databases are also incorporated by reference.

## EXAMPLES

### EXAMPLE 1

#### Generation of Plants with a HIO32.3 Phenotype by Transformation with an Activation Tagging Construct

[0057] Mutants were generated using the activation tagging "ACTTAG" vector, pSKI015 (GI#6537289; Weigel D *et al.*, 2000). Standard methods were used for the generation of *Arabidopsis* transgenic plants, and were essentially as described in published application PCT WO0183697. Briefly, T0 *Arabidopsis* (Col-0) plants were transformed with *Agrobacterium* carrying the pSKI015 vector, which comprises T-DNA derived from the *Agrobacterium* Ti plasmid, an herbicide resistance selectable marker gene, and the 4X CaMV 35S enhancer element. Transgenic plants were selected at the T1 generation based on herbicide resistance. T2 seed was collected from T1 plants and stored in an indexed collection, and a portion of the T2 seed was accessed for the screen.

[0058] Quantitative determination of seed fatty acid content was performed using the following methods. A sample of 15 to 20 T2 seeds from each line tested, which generally contained homozygous insertion, homozygous wild-type, and heterozygous genotypes in a standard 1:1:2 ratio, was massed on UMT-2 ultra-microbalance (Mettler-Toledo Co., Ohio, USA) and then transferred to a glass extraction vial. Lipids were extracted from the seeds and trans-esterified in 500 ul 2.5% H<sub>2</sub>SO<sub>4</sub> in MeOH for 3 hours at 80 °C, following the method of Browse et al. (Biochem J 235:25-31, 1986) with modifications. A known amount of heptadecanoic acid was included in the reaction as an internal standard. 750 ul of water and 400 ul of hexane were added to each vial, which was then shaken vigorously and allowed to phase separate. Reaction vials were loaded directly onto GC for analysis and the upper hexane phase was sampled by the autosampler. Gas chromatography with Flame Ionization detection was used to separate and quantify the fatty acid methyl esters. Agilent 6890 Plus GC's were used for separation with Agilent Innowax columns (30m x 0.25mm ID, 250um film thickness). The carrier gas was Hydrogen at a constant flow of 2.5 ml/ minute. 1ul of sample was injected in splitless mode (inlet temperature 220°C, Purge flow 15ml/min at 1 minute). The oven was programmed for an initial temperature of 105°C, initial time 0.5 minutes, followed by a ramp of 60°C per minute to 175°C, a 40°C /minute ramp to 260°C with a final hold time of 2 minutes. Detection was by Flame Ionization (Temperature 275°C, Fuel flow 30.0 ml/min, Oxidizer 400.0 ml/min). Instrument control and data collection and analysis were monitored using the Millennium Chromatography Management System (Version 3.2, Waters Corporation, Milford, MA). Integration and quantification were performed automatically, but all analyses were subsequently examined manually to verify correct peak identification and acceptable signal to noise ratio before inclusion of the derived results in the study.

[0059] The ACTTAG line designated W000082263 was identified as having a high oil phenotype. Specifically, oil constituted 36.3% of seed mass (w/w) compared to an average oil content of 27.0% of other ACTTAG lines grown and analyzed in the same conditions (e.g., reference lines). Reanalysis of the same seed was performed in duplicate. This analysis confirmed an increase in oil content relative to controls. It was concluded that the presence of the ACTTAG locus can increase seed oil content between 7% and 11% relative to controls. It is determined that the high oil phenotype is dominant based on oil content of seeds from genotyped individuals.

## **EXAMPLE 2**

### **Characterization of the T-DNA Insertion in Plants Exhibiting the Altered Oil Content Phenotype.**

[0060] We performed standard molecular analyses, essentially as described in patent application PCT WO0183697, to determine the site of the T-DNA insertion associated

with the altered oil content phenotype. Briefly, genomic DNA was extracted from plants exhibiting the altered oil content phenotype. PCR, using primers specific to the pSKI015 vector, confirmed the presence of the 35S enhancer in plants from line W000082263, and Southern blot analysis verified the genomic integration of the ACTTAG T-DNA and showed the presence of a single T-DNA insertion in the transgenic line.

[0061] Plasmid rescue was used to recover genomic DNA flanking the T-DNA insertion, which was then subjected to sequence analysis using a basic BLASTN search of the *Arabidopsis* Genomic DNA (TAIR) database (see publicly available Arabidopsis Information Resource website). There was sequence identity to BAC clone T23J7 (GI#4741184), chromosome 3. Sequences from nucleotides 16030-16423 and 16443-17200 were recovered, placing the left border junction upstream of nucleotide 16423 (GI#4741184) and downstream of nucleotide 15443. The sequences from nucleotides 15424-16442 of BAC clone T23J7, chromosome 3 (GI#4741184) are deleted in the mutant chromosome.

[0062] To determine whether the T-DNA insertion causes the high seed oil phenotype cosegregation of the high seed oil phenotype and the presence of the T-DNA was tested. Eighteen T2 plants were grown to maturity and seed harvested from these plants was used to determine the seed oil phenotype. The seed oil content from these was determined as described in Example 1. The genotype of the T2 seed was inferred by analyzing the T3 seed for the presence or absence of the T-DNA at the site of the insertion by PCR using primers that are specific to the corresponding genomic region and the T-DNA. The results show that the average oil content of T3 seed containing the T-DNA insert was higher than those families lacking the insert. T2 individuals homozygous for the T-DNA at this locus produced seed with an oil content of 107.5% compared to progeny lacking the T-DNA. T2 individuals hemizygous for these loci produced seed with an oil content of 112.7% compared to progeny lacking the T-DNA. Because the homozygotes and hemizygotes for the high oil loci display a similar increase in oil content, we conclude that the T-DNA is linked with the high oil phenotype and the phenotype is caused by a dominant mutation.

[0063] Sequence analysis revealed that the left border of the T-DNA insert was about 494 base pairs 5' of the translation start site of the gene whose nucleotide sequence is presented as SEQ ID NO: 1 which we designated HIO32.3.

**EXAMPLE 3****Analysis of *Arabidopsis* HIO32.3 Sequence**

[0064] Sequence analyses were performed with BLAST (Altschul *et al.*, 1990, J. Mol. Biol. 215:403-410), PFAM (Bateman *et al.*, 1999, Nucleic Acids Res 27:260-262), PSORT (Nakai K, and Horton P, 1999, Trends Biochem Sci 24:34-6), and/or CLUSTAL (Thompson JD *et al.*, 1994, Nucleic Acids Res 22:4673-4680).

BLASTN against ESTs:

[0065] There are more than 10 *Arabidopsis* ESTs (unigene cluster At.9308) that correspond to the *Arabidopsis* paralogue At5g62520. No EST corresponds to the candidate gene At3g47720 was identified.

[0066] There are also other similar plant ESTs showing similarity to At3g47720. If possible, ESTs contigs of each species were made. The best hit for each of the following species are listed below and included in the "Orthologue Table" below: *Brassica rapa*, *Glycine max*, *Lycopersicon esculentum*, *Gossypium hirsutum*, *Zea mays*, *Oryza sativa*, *Triticum aestivum*, *Helianthus annuus*, *Solanum tuberosum*.

1. One EST from Chinese cabbage (*Brassica rapa* subsp. *pekinensis*)  
>gi|20374907|gb|BG543927.1|BG543927 E1677 Chinese cabbage etiolated seedling library  
Score = 329 (120.9 bits), Expect = 2.7e-35

2. One EST contig from soybean  
>gi|19269392|gb|BM885648.1|BM885648 from germinating shoot (24 hr germination)  
>gi|15285085|gb|BI468976.1|BI468976 from 3 week old whole seedlings  
Score = 318 (117.0 bits), Expect = 4.0e-34

The contigged sequence is provided below as SEQ ID NO:3.

3. One EST from cotton  
>gi|3325723|gb|AI054609.1|AI054609 coau0001I06 Cotton Boll Abscission Zone cDNA library  
Score = 223 (83.6 bits), Expect = 4.7e-24

4. One EST contig from maize  
>gi|14203679|gb|BG837356.1|BG837356 Zm10\_08c08\_A ESTs from Maize Silk Six Hours After Silk Channel Inoculation with *Fusarium graminearum*  
>gi|4646391|gb|AI621721.1|AI621721 486006G12.x5 486 - leaf primordia cDNA library from Hake lab  
>gi|4217842|gb|AI391838.1|AI391838 486006A11.x1 486 - leaf primordia cDNA library from Hake lab  
>gi|4217921|gb|AI391917.1|AI391917 486005B03.x1 486 - leaf primordia cDNA library from Hake lab  
Score = 176 (67.0 bits), Expect = 1.8e-26

The contigged sequence is provided below as SEQ ID NO:4.

5. One EST from sunflower (*Helianthus annuus*)

>gi|22387461|gb|BQ969940.1|BQ969940 QHB39P16.yg.ab1 QH\_ABCDI sunflower RHA801  
isolated from a library constructed with different parts of the plant  
Score = 179 (68.1 bits), Expect = 5.7e-30

Note that the following ESTs show significant homology to >gi|22387461. It is possible that these sequences represent of the same mRNA and that the differences in nucleotide sequences may be the result of sequencing error or of single nucleotide polymorphisms.

>gi|22387167|gb|BQ969646.1|BQ969646  
>gi|22315760|gb|BQ916979.1|BQ916979  
>gi|22315563|gb|BQ916782.1|BQ916782  
>gi|22313791|gb|BQ915010.1|BQ915010  
>gi|22315308|gb|BQ916527.1|BQ916527  
>gi|22387224|gb|BQ969703.1|BQ969703  
>gi|22388939|gb|BQ971418.1|BQ971418  
>gi|22389044|gb|BQ971523.1|BQ971523  
>gi|22386431|gb|BQ968910.1|BQ968910

5. One EST from tomato (*Lycopersicon esculentum*)

>gi|4387847|gb|AI483923.1|AI483923 EST249794 tomato ovary, TAMU *Lycopersicon*  
*esculentum*  
Score = 241 (89.9 bits), Expect = 5.8e-26

6. One EST from wheat (*Triticum aestivum*)

>gi|24979906|gb|CA485901.1|CA485901 WHE4324\_E06\_J12ZS Wheat meiotic  
anther cDNA  
Score = 186 (70.5 bits), Expect = 9.1e-30

The EST described above (>gi|24979906) encompasses the sequence from the following EST. However, these two sequences differ by one nucleotide that could be the result of sequencing error or of single nucleotide polymorphism

>gi|24978183|gb|CA484178.1|CA484178 WHE4303\_C01\_F01ZS Wheat meiotic anther  
cDNA

7. One EST from potato (*Solanum tuberosum*)

>gi|13616454|gb|BG598314.1|BG598314 EST496992 cSTS *Solanum tuberosum* cDNA clone  
cSTS20P7 5' sequence, isolated from sprouting eyes of tubers  
Score = 191 (72.3 bits), Expect = 1.8e-28

8. One EST from rice (*Oryza sativa*)

>gi|18385875|gb|BM419074.1|BM419074 R009A04 *Oryza sativa* mature leaf library induced  
by *M. grisea*  
Score = 153 (58.9 bits), Expect = 4.3e-16

Note that the following ESTs show significant homology to >gi|18385875. It is possible that these sequences represent of the same mRNA and that the differences in nucleotide sequences may be the result of sequencing error or of single nucleotide polymorphisms.

>gi|22307912|gb|BQ909134.1|BQ909134  
>gi|3762808|dbj|AU029560.1|AU029560

## BLASTP against amino acids:

[0067] SEQ ID NO:2 has a high degree of homology to two plant proteins and low degree of homology to a human protein. The top 7 BLAST hits are listed and are included in the "Orthologue Table" below.

## 1. Itself (3 redundant entries)

>gi|15228235|ref|NP\_190356.1| putative protein; protein id: At3g47720.1 [Arabidopsis thaliana]

>gi|7487375|pir|T07711 hypothetical protein T23J7.50 - Arabidopsis thaliana

>gi|4741189|emb|CAB41855.1| putative protein [Arabidopsis thaliana]

Score = 604 bits (1557), Expect = e-172

## 2. At5g62520 from Arabidopsis (several redundant entries)

>gi|15241862|ref|NP\_201058.1| putative protein; protein id: At5g62520.1, supported by cDNA: 23920., supported by cDNA: gi\_15450638, supported by cDNA: gi\_17386159 [Arabidopsis thaliana]

>gi|10178083|dbj|BAB11502.1| gene\_id:K19B1.13~pir|T07711~similar to unknown protein [Arabidopsis thaliana]

>gi|15450639|gb|AAK96591.1| AT5g62520/K19B1\_13 [Arabidopsis thaliana]

>gi|17386160|gb|AAL38626.1|AF446893\_1 AT5g62520/K19B1\_13 [Arabidopsis thaliana]

Score = 281 bits (718), Expect = 1e-74

The following sequence is highly likely to be a redundant entry of At5g62520 because tblastn results indicate that they are localized to the same place in the Arabidopsis genome. However, it differs from the sequences listed above by two amino acids. This is likely to be the result of sequencing errors or of single nucleotide polymorphisms with little or no effect on activity.

>gi|21592445|gb|AAM64396.1| unknown [Arabidopsis thaliana]

Score = 275 bits (704), Expect = 5e-73

## 3. At1g23550 from Arabidopsis (3 redundant entries)

>gi|15220787|ref|NP\_173769.1| hypothetical protein; protein id: At1g23550.1 [Arabidopsis thaliana]

>gi|4056438|gb|AAC98011.1| F5O8.11 [Arabidopsis thaliana]

Score = 158 bits (399), Expect = 1e-37

The following sequence is another redundant entry of At1g23550 based on Genbank annotation. However, it differs from the sequences listed above by one amino acid. This is likely to be the result of sequencing errors or of single nucleotide polymorphisms with little or no effect on activity.

>gi|8778582|gb|AAF79590.1|AC007945\_10 F28C11.18 [Arabidopsis thaliana]

Score = 154 bits (390), Expect = 1e-36

## 4. At1g70440 from Arabidopsis (3 redundant entries)

>gi|15223159|ref|NP\_177201.1| hypothetical protein; protein id: At1g70440.1 [Arabidopsis thaliana]

Score = 131 bits (330), Expect = 1e-29

The following sequences are redundant entries of At1g70440 based on Genbank annotation. However, they differ from the sequence listed above by several amino acids. This is likely to be the result of sequencing errors or of single nucleotide polymorphisms with little or no effect on activity.

>gi|7485680|pir|T01478 hypothetical protein F17O7.2 - Arabidopsis thaliana

>gi|3176692|gb|AAC18815.1| F17O7.2 [Arabidopsis thaliana]

Score = 118 bits (296), Expect = 8e-26

5. At1g32230 from Arabidopsis (several redundant entries)

>gi|18398335|ref|NP\_564391.1| expressed protein; protein id: At1g32230.1, supported by cDNA: gi\_11044956, supported by cDNA: gi\_19715642 [Arabidopsis thaliana]

>gi|19715643|gb|AAL91641.1| At1g32230/F3C3\_1 [Arabidopsis thaliana]

Score = 111 bits (278), Expect = 1e-23

The following sequences are highly likely to be a redundant entry of At5g62520 based on (1) tblastn results indicate that they are localized to the same place in the Arabidopsis genome, and (2) Genbank annotation. However, they differ from the sequences listed above by a few amino acids. This is likely to be the result of sequencing errors or of single nucleotide polymorphisms with little or no effect on activity.

>gi|16604703|gb|AAL24144.1| unknown protein [Arabidopsis thaliana]

>gi|23296877|gb|AAN13193.1| unknown protein [Arabidopsis thaliana]

>gi|11044957|emb|CAC14428.1| ceo protein [Arabidopsis thaliana]

>gi|10801372|gb|AAG23444.1|AC084165\_10 unknown protein [Arabidopsis thaliana]

>gi|14150488|gb|AAK54509.1|AF317898\_1 ATP8 [Arabidopsis thaliana]

6. A putative CEO protein from rice

>gi|18057156|gb|AAL58179.1|AC027037\_1 putative CEO protein [Oryza sativa]

Score = 99.4 bits (246), Expect = 7e-20

7. At2g35510 from Arabidopsis (several redundant entries)

>gi|18403862|ref|NP\_565806.1| expressed protein; protein id: At2g35510.1, supported by cDNA: gi\_15810142 [Arabidopsis thaliana]

>gi|20197351|gb|AAC36170.2| expressed protein [Arabidopsis thaliana]

>gi|23297349|gb|AAN12947.1| unknown protein [Arabidopsis thaliana]

Score = 85.5 bits (210), Expect = 9e-16

[0068] The following sequence is another redundant entry of At1g23550 based on Genbank annotation. However, it differs from the sequences listed above by one amino acid. This is likely to be the result of sequencing errors or of single nucleotide polymorphisms with little or no effect on activity.

>gi|15810143|gb|AAL07215.1| unknown protein [Arabidopsis thaliana]

Score = 84.0 bits (206), Expect = 3e-15

Ortholog Gene Name	Species	GI #	% ID to HIO32.3	Score(s) (BLAST, Clustal, etc.)	Coordinate s of protein motif (s)	%ID to HIO32.3 motif or to Pfam/other consensus
A putative CEO protein from rice	Oryza sativa	>gi 18057156 56	Length = 591 Identities = 67/256 (26%), Positives = 120/256 (46%)	BLASTP Score = 99.4 bits (246), Expect = 7e-20		
A Chinese cabbage EST	Brassica rapa subsp. pekinensis	>gi 20374907	Partial sequenc, using reading frame +2 Length = 198 Identities = 86/195 (44%), Positives = 118/195 (60%)	TBLASTN Score = 329 (120.9 bits), Expect = 2.7e-35, P = 2.7e-35		

An EST contig from soybean	Glycine max	92 85 >gi 192693 >gi 152850	Partial sequenc, using reading frame +1 Length = 256 Identities = 85/230 (36%), Positives = 128/230 (55%)	TBLASTN Score = 318 (117.0 bits), Expect = 4.0e-34, P = 4.0e-34		
An EST from cotton boll abscission zone	Gossypiu m hirsutum	3 >gi 332572	Partial sequenc, using reading frame +2 Length = 79 Identities = 40/72 (55%), Positives = 56/72 (77%)	TBLASTN Score = 223 (83.6 bits), Expect = 4.7e-24, P = 4.7e-24		
An EST from maize	Zea mays	79 1 2 1 >gi 142036 >gi 464639 >gi 421784 >gi 421792	Partial sequenc, using reading frame -1 Length = 344 Identities = 55/209 (26%), Positives = 88/209 (42%)	TBLASTN Score = 176 (67.0 bits), Expect = 1.8e-26, Sum P(2) = 1.8e-26		
A sunflower EST	Helianthu s annuus	61 >gi 223874	Partial sequenc, using reading frame +1 Length = 232 Identities = 37/88 (42%), Positives = 54/88 (61%)	TBLASTN Score = 179 (68.1 bits), Expect = 5.7e-30, Sum P(2) = 5.7e-30		
A tomato ovary EST	Lycopers icon esculentum	7 >gi 438784	Partial sequenc, using reading frame Length = 135 Identities = 49/106 (46%), Positives = 70/106 (66%)	TBLASTN Score = 241 (89.9 bits), Expect = 5.8e-26, P = 5.8e-26		
A wheat EST	Triticum aestivum	06 >gi 249799	Partial sequenc, using reading frame +1 Length = 230 Identities = 40/94 (42%), Positives = 56/94 (59%)	TBLASTN Score = 186 (70.5 bits), Expect = 9.1e-30, Sum P(2) = 9.1e-30		
A potato EST	Solanum tuberosum	54 >gi 136164	Partial sequenc, using reading frame +1 Length = 189 Identities = 31/71 (43%), Positives = 53/71 (74%)	TBLASTN Score = 191 (72.3 bits), Expect = 1.8e-28, Sum P(2) = 1.8e-28		
A rice EST	Oryza sativa	75 >gi 183858	Partial sequenc, using reading frame +3 Length = 122 Identities = 29/69 (42%), Positives = 48/69 (69%)	TBLASTN Score = 153 (58.9 bits), Expect = 4.3e-16, P = 4.3e-16		



**Closest Arabidopsis homologs:**

At5g62520	Arabidopsis thaliana	62 >gi 152418 83 >gi 101780 39 >gi 154506 60 >gi 173861	Length = 309 Identities = 157/301 (52%), Positives = 200/301 (66%)	BLASTP Score = 281 bits (718), Expect = 1e-74		
At1g23550	Arabidopsis thaliana	87 >gi 152207 8 >gi 405643	Length = 323 Identities = 90/261 (34%), Positives = 152/261 (58%)	BLASTP Score = 158 bits (399), Expect = 1e-37		
At1g70440	Arabidopsis thaliana	59 >gi 152231	Length = 305 Identities = 76/240 (31%), Positives = 138/240 (57%)	BLASTP Score = 131 bits (330), Expect = 1e-29		
At1g32230	Arabidopsis thaliana	35 >gi 183983 43 >gi 197156	Length = 589 Identities = 74/284 (26%), Positives = 127/284 (44%)	BLASTP Score = 111 bits (278), Expect = 1e-23	PF02825: aa 78-149	PF02825 : Score = 42.9, E value = 7.6e-10
At2g35510	Arabidopsis thaliana	62 >gi 184038 51 >gi 201973 49 >gi 232973	Length = 568 Identities = 75/284 (26%), Positives = 120/284 (42%)	BLASTP Score = 85.5 bits (210), Expect = 9e-16	PF02825: aa 77-148	PF02825 : Score = 80.3, E value = 4.4e-21

[0069] At3g47720 is a non-secretory protein, lacks signal peptide (predicted by signalP) and has no transmembrane domain (predicted by TMHMM). The cellular localization of At3g47720 remains unclear based on PSORT2 analysis (48% cytoplasmic, 32% nuclear, 20% cytoskeletal). No known protein motif can be detected for At3g47720 by Pfam analysis.

[0070] At3g47720 encodes a protein that has not been functionally characterized and has no known protein motif. BLAST results of At3g47720 suggest that it is a member of small protein family unique to plants.

#### **EXAMPLE 4**

##### **Confirmation of Phenotype/Genotype Association**

[0071] RT-PCR analysis showed that the HIO32.3 gene was overexpressed in plants from the line displaying the HIO32.3 phenotype. Specifically, RNA was extracted from rosette leaves and/or siliques of plants exhibiting the HIO32.3 phenotype collected at a variety of developmental stages and pooled. RT-PCR was performed using primers specific to the

sequence presented as SEQ ID NO:1, to other predicted genes in the vicinity of the T-DNA insertion, and to a constitutively expressed actin gene (positive control). The results showed that in plants displaying the HIO32.3 phenotype, mRNA for the HIO32.3 gene is up-regulated in leaves.

[0072] The dominant inheritance pattern of the HIO32.3 phenotype is confirmed through genetic analysis. In general, genetic analysis involves the production and analysis of F1 hybrids. Typically, F1 crosses are carried out by collecting pollen from T2 plants, which is used to pollinate wild type plants. Such crosses are carried out by taking about 4 flowers from each selected individual plants, and using the T2 flower as the male pollen donor and flowers of the wild type plants as the female. 4-5 crosses are done for an individual of interest. Seed formed from crosses of the same individual are pooled, planted and grown to maturity as F1 hybrids.

#### **EXAMPLE 5**

##### **Recapitulation of the High Oil Phenotype**

[0073] To confirm whether over-expression of At3g47720 causes a high seed oil phenotype, oil content in seeds from transgenic plants over-expressing this gene was compared with oil content in seeds from non-transgenic control plants. To do this, At3g47720 was cloned into a plant transformation vector behind the strong constitutive CsVMV promoter and transformed into Arabidopsis plants using the floral dip method. The plant transformation vector contains the nptII gene, which provides resistance to kanamycin, and serves as a selectable marker. Seed from the transformed plants were plated on agar medium containing kanamycin. After 7 days, transgenic plants were identified as healthy green plants and transplanted to soil. Non-transgenic control plants were germinated on agar medium, allowed to grow for 7 days and then transplanted to soil. Twenty-two transgenic seedlings and 10 non-transgenic control plants were transplanted to random positions in the same 32 cell flat. The plants were grown to maturity, allowed to self-fertilize and set seed. Seed was harvested from each plant and its oil content estimated by Near Infrared (NIR) Spectroscopy using methods described below.

[0074] NIR infrared spectra were captured using a Bruker 22 N/F near infrared spectrometer. Bruker Software was used to estimate total seed oil and total seed protein

content using NIR data from the samples and reference methods according to the manufacturer's instructions. An oil content predicting calibration was developed following the general method of AOCS Procedure Am1-92, Official Methods and Recommended Practices of the American Oil Chemists Society, 5th Ed., AOCS, Champaign Ill). The calibration allowing NIR predictions of Crude Oil Crude Oil ASE (Ren Oil, Accelerated Solvent Extraction) was developed.

[0075] The effect of over-expression of At3g47720 on seed oil has been tested in five experiments. In three experiments, the plants over-expressing At3g47720 had higher seed oil content than the control plants grown in the same flat; there was no difference in two experiments. Across the experiments, the average seed oil content of plants over-expressing At3g47720 was 3.8% greater than the untransformed controls. The in seed oil content in plants over-expressing At3g47720 was significantly greater than non-transgenic control plants (two-way ANOVA;  $P < 0.0001$ ).

Experiment	Plant ID	Transgene	Predicted average	Relative value average
1	Z003905001	CsVMV::HIO32.3	34.1198	105.0453
1	Z003905002	CsVMV::HIO32.3	34.6113	106.5584
1	Z003905003	CsVMV::HIO32.3	34.9244	107.5223
1	Z003905004	CsVMV::HIO32.3	32.3115	99.4779
1	Z003905005	CsVMV::HIO32.3	34.6011	106.5269
1	Z003905006	CsVMV::HIO32.3	35.4656	109.1885
1	Z003905007	CsVMV::HIO32.3	35.5339	109.3987
1	Z003905008	CsVMV::HIO32.3	32.6891	100.6404
1	Z003905009	CsVMV::HIO32.3	31.1428	95.8799
1	Z003905010	CsVMV::HIO32.3	33.3398	102.6437
1	Z003905011	CsVMV::HIO32.3	34.0836	104.9338
1	Z003905012	CsVMV::HIO32.3	33.8851	104.3226
1	Z003905013	CsVMV::HIO32.3	32.6806	100.6142
1	Z003905014	CsVMV::HIO32.3	35.4043	108.9999
1	Z003905015	CsVMV::HIO32.3	32.4204	99.8133
1	Z003905016	CsVMV::HIO32.3	34.2276	105.3769
1	Z003905017	CsVMV::HIO32.3	35.9069	110.5472
1	Z003905018	CsVMV::HIO32.3	35.1153	108.1099
1	Z003905019	CsVMV::HIO32.3	33.4684	103.0396
1	Z003905020	CsVMV::HIO32.3	36.723	113.0598
1	Z003905021	CsVMV::HIO32.3	33.5667	103.3424
1	Z003905022	CsVMV::HIO32.3	34.2919	105.5749
1	Z003895001	None	34.0726	104.8999
1	Z003895002	None	34.0624	104.8683

1	Z003895003	None	32.1291	98.9163
1	Z003895004	None	30.4016	93.5978
1	Z003895005	None	33.8228	104.1308
1	Z003895006	None	32.6772	100.604
1	Z003895007	None	31.917	98.2634
1	Z003895008	None	32.2821	99.3876
1	Z003895009	None	32.1418	98.9556
1	Z003895010	None	31.3041	96.3764
2	Z003983001	CsVMV::HIO32.3	34.4143	99.8947
2	Z003983002	CsVMV::HIO32.3	35.1919	102.1518
2	Z003983003	CsVMV::HIO32.3	35.3644	102.6526
2	Z003983004	CsVMV::HIO32.3	35.1895	102.1449
2	Z003983005	CsVMV::HIO32.3	31.8284	92.3886
2	Z003983006	CsVMV::HIO32.3	34.9055	101.3204
2	Z003983007	CsVMV::HIO32.3	32.4582	94.2168
2	Z003983008	CsVMV::HIO32.3	34.2324	99.3666
2	Z003983009	CsVMV::HIO32.3	31.8428	92.4302
2	Z003983010	CsVMV::HIO32.3	36.2317	105.1699
2	Z003983011	CsVMV::HIO32.3	36.096	104.7762
2	Z003983012	CsVMV::HIO32.3	34.7108	100.7555
2	Z003983013	CsVMV::HIO32.3	33.6519	97.6817
2	Z003983014	CsVMV::HIO32.3	35.2501	102.3209
2	Z003983016	CsVMV::HIO32.3	34.9047	101.318
2	Z003983017	CsVMV::HIO32.3	35.3804	102.6991
2	Z003983018	CsVMV::HIO32.3	33.7934	98.0923
2	Z003983019	CsVMV::HIO32.3	30.4627	88.4243
2	Z003983020	CsVMV::HIO32.3	34.5856	100.3919
2	Z003983021	CsVMV::HIO32.3	36.5568	106.1138
2	Z003983022	CsVMV::HIO32.3	37.2214	108.0428
2	Z003999001	None	34.4392	99.967
2	Z003999002	None	35.3728	102.6768
2	Z003999003	None	34.9726	101.5152
2	Z003999004	None	34.8285	101.0971
2	Z003999005	None	34.3824	99.8022
2	Z003999006	None	32.4696	94.2499
2	Z003999007	None	36.503	105.9575
2	Z003999008	None	34.9916	101.5703
2	Z003999009	None	32.8391	95.3225
2	Z003999010	None	33.707	97.8415
3	DX02966001	CsVMV::HIO32.3	30.8834	89.4963
3	DX02966002	CsVMV::HIO32.3	33.0238	95.6992
3	DX02966003	CsVMV::HIO32.3	32.1994	93.3102
3	DX02966004	CsVMV::HIO32.3	34.847	100.9824
3	DX02966005	CsVMV::HIO32.3	33.0958	95.9076
3	DX02966006	CsVMV::HIO32.3	34.4815	99.9234
3	DX02966007	CsVMV::HIO32.3	33.8108	97.9797
3	DX02966008	CsVMV::HIO32.3	35.14	101.8317
3	DX02966009	CsVMV::HIO32.3	35.7125	103.4906

3	DX02966010	CsVMV::HIO32.3	34.0653	98.7171
3	DX02966011	CsVMV::HIO32.3	34.7922	100.8238
3	DX02966012	CsVMV::HIO32.3	36.2254	104.9768
3	DX02966013	CsVMV::HIO32.3	35.6025	103.1719
3	DX02966015	CsVMV::HIO32.3	36.3922	105.4602
3	DX02965001	None	34.3094	99.4246
3	DX02965002	None	33.8394	98.0626
3	DX02965003	None	34.7631	100.7395
3	DX02965004	None	35.2757	102.2248
3	DX02965005	None	34.4162	99.734
3	DX02965006	None	34.8555	101.0071
3	DX02965007	None	33.7694	97.8598
3	DX02965008	None	34.8349	100.9476
4	DX06939001	CsVMV::HIO32.3	33.2057	113.2141
4	DX06939002	CsVMV::HIO32.3	30.9565	105.5454
4	DX06939003	CsVMV::HIO32.3	30.7753	104.9276
4	DX06939004	CsVMV::HIO32.3	32.2312	109.8916
4	DX06939005	CsVMV::HIO32.3	34.8463	118.8074
4	DX06939006	CsVMV::HIO32.3	32.041	109.2431
4	DX06939007	CsVMV::HIO32.3	34.7717	118.5533
4	DX06939008	CsVMV::HIO32.3	32.7271	111.5822
4	DX06939009	CsVMV::HIO32.3	30.5371	104.1153
4	DX06939010	CsVMV::HIO32.3	28.8958	98.5195
4	DX06939011	CsVMV::HIO32.3	31.6481	107.9033
4	DX06939012	CsVMV::HIO32.3	36.3548	123.9507
4	DX06939013	CsVMV::HIO32.3	34.8001	118.6501
4	DX06939014	CsVMV::HIO32.3	31.4454	107.2121
4	DX06939015	CsVMV::HIO32.3	31.9814	109.0399
4	DX06939016	CsVMV::HIO32.3	31.8235	108.5015
4	DX06939017	CsVMV::HIO32.3	30.9507	105.5257
4	DX06939018	CsVMV::HIO32.3	30.9881	105.6532
4	DX06939020	CsVMV::HIO32.3	32.0067	109.1261
4	DX06939021	CsVMV::HIO32.3	35.2219	120.0881
4	DX06939022	CsVMV::HIO32.3	32.503	110.8181
4	DX06957001	None	28.8224	98.2691
4	DX06957002	None	29.3474	100.0591
4	DX06957003	None	28.7321	97.9615
4	DX06957004	None	30.1454	102.78
4	DX06957005	None	32.6027	111.1581
4	DX06957006	None	28.8494	98.3612
4	DX06957007	None	27.4123	93.4616
4	DX06957008	None	30.1262	102.7144
4	DX06957009	None	27.6125	94.1441
4	DX06957010	None	29.65	101.0908
5	DX06940001	CsVMV::HIO32.3	28.7458	97.9233
5	DX06940002	CsVMV::HIO32.3	27.5702	93.9185
5	DX06940003	CsVMV::HIO32.3	30.1592	102.7378
5	DX06940004	CsVMV::HIO32.3	30.5856	104.1905

5	DX06940005	CsVMV::HIO32.3	28.7915	98.079
5	DX06940006	CsVMV::HIO32.3	29.363	100.0258
5	DX06940007	CsVMV::HIO32.3	31.1195	106.0091
5	DX06940008	CsVMV::HIO32.3	29.2426	99.6156
5	DX06940009	CsVMV::HIO32.3	31.056	105.7928
5	DX06940010	CsVMV::HIO32.3	30.3929	103.5341
5	DX06940011	CsVMV::HIO32.3	29.53	100.5944
5	DX06940012	CsVMV::HIO32.3	33.4601	113.9826
5	DX06940013	CsVMV::HIO32.3	29.4551	100.3392
5	DX06940014	CsVMV::HIO32.3	28.9251	98.5341
5	DX06940015	CsVMV::HIO32.3	30.0224	102.2717
5	DX06940016	CsVMV::HIO32.3	28.6861	97.7199
5	DX06940017	CsVMV::HIO32.3	33.3423	113.5813
5	DX06940018	CsVMV::HIO32.3	31.4592	107.1666
5	DX06940019	CsVMV::HIO32.3	29.8827	101.7959
5	DX06940020	CsVMV::HIO32.3	31.4644	107.1843
5	DX06940021	CsVMV::HIO32.3	28.3517	96.5805
5	DX06940022	CsVMV::HIO32.3	31.4847	107.2534
5	DX06958001	None	29.3962	100.1388
5	DX06958002	None	29.9035	101.867
5	DX06958003	None	30.7418	104.7225
5	DX06958004	None	29.5126	100.5353
5	DX06958005	None	27.8422	94.845
5	DX06958006	None	27.849	94.8681
5	DX06958007	None	29.8981	101.8484
5	DX06958008	None	29.6368	100.9582
5	DX06958009	None	29.5021	100.4996
5	DX06958010	None	29.2724	99.7171

**REFERENCES**

- Altschul, S.F. *et al.*, *J. Mol. Biol.* 215:403-410, 1990.
- Altschul, S.F. *et al.*, *Nucleic Acids Res.* 25:3389-3402, 1997.
- Ausubel FM *et al.* Current Protocols in Molecular Biology, John Wiley & Sons, New York, N.Y., 1993.
- Baldwin D *et al.*, *Cur Opin Plant Biol.* 2(2):96-103, 1999.
- Bateman *et al.*, 1999, *Nucleic Acids Res* 27:260-262.
- Baulcombe D, *Arch Virol Suppl* 15:189-201, 1999.
- Cannon *et al.*, *Plant Molec. Biol.* (1990) 15:39-47.
- Ch'ng *et al.*, *Proc. Natl. Acad. Sci. USA* (1989) 86:10006-10010
- Christensen S *et al.*, 9<sup>th</sup> International Conference on *Arabidopsis* Research. Univ. of Wisconsin-Madison, June 24-28, 1998. Abstract 165.
- Christou *et al.*, *Proc. Natl. Acad. Sci USA* (1989) 86:7500-7504.
- Cough, SJ and Bent, AF, *the Plant Journal* 16(6): 735-743, 1998.
- De Block *et al.*, *Plant Physiol.* (1989) 91:694-701.
- Dieffenbach C and Dveksler G (Eds.) PCR Primer: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY, 1989.
- Everett *et al.*, *Bio/Technology* (1987) 5:1201
- Feldmann *et al.*, *Science* 243: 1351-1354, 1989.
- Focks N and Benning C, *Plant Physiol* 118:91-101, 1998.
- Fridborg I *et al.*, *Plant Cell* 11: 1019-1032, 1999.
- Geest AH and Hall TC, *Plant Mol Biol* 32(4):579-88, 1996.
- Gelvin, S. B., Schilperoort, R. A., Varma, D. P. S., eds. *Plant Molecular Biology Manual* 1990.
- Glick, BR and Thompson, JE, Eds. *METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY*, p. 213-221, CRC Press, 1993.
- Harlow E and Lane D, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1988, New York.
- Harlow E and Lane D, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1999, New York
- Hayashi H *et al.*, *Science* 258: 1350-1353, 1992.
- James DW and Dooner HK (1990) *Theor Appl Genet* 80, 241-245.
- Jensen, L.G., *et al.*, *Proc. Natl. Acad. Sci. USA* 93:3487-3491, 1996.
- Jones JD *et al.*, *Transgenic Res* 1:285-297 1992.

- Kardailsky I *et al.*, *Science* 286: 1962-1965, 1999.
- Kline *et al.*, *Nature* (1987) 327:70.
- Kunkel TA *et al.*, *Methods Enzymol.* 204:125-39, 1991.
- Lemieux B, *et al.* 1990, *Theor Appl Genet* 80, 234-240.
- Nakamura Y *et al.*, 1999, *Nucleic Acids Res* 27:292.
- Napoli, *et al.*, *Plant Cell* 2:279-289, 1990.
- Omirulleh *et al.*, *Plant Mol Biol.* 21(3):415-28, 1993.
- Sambrook *et al.* *Molecular Cloning: A Laboratory Manual* (Second Edition), Cold Spring Harbor Press, Plainview, N.Y., 1989.
- Schaffer R, *et al.*, *Cell* 93: 1219-1229, 1998.
- Sheehy *et al.*, *Proc. Natl. Acad. Sci. USA* (1988) 85:8805-8809.
- Smith, *et al.*, *Nature* 334:724-726, 1988.
- Smith *et al.*, *Mol. Gen. Genetics* (1990) 224:477-481.
- Thompson JD *et al.*, 1994, *Nucleic Acids Res* 22:4673-4680.
- van der Krol *et al.*, *Biotechniques* (1988) 6:958-976.
- van der Krol *et al.*, *The Plant Cell* (1990) 2:291-299.
- Van Haaren MJJ *et al.*, *Plant Mol Bio* 21:625-640, 1993.
- Verdaguer B *et al.*, *Plant Mol Biol* 37:1055-1067, 1998.
- Waterhouse, *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964, 1998.
- Weigel D, *et al.*, *Plant Physiology*, 122:1003-1013, 2000.
- Wilson K *et al.*, *Plant Cell* 8: 659-671, 1996.
- Yadav NS *et al.* (1993) *Plant Physiol* 103, 467-476.
- Bert *et al.*, *Theor Appl Genet.* 2003 Jun;107(1):181-9.
- Colbert *et al.* (2001) *Plant Physiol* 126:480-484.
- Jako, *et al.*, *Plant Physiology* 126(2):861-74, 2001.
- Katavic, *et al.*, *Plant Physiology*, 108(1):399-409, 1995.
- Lionneton *et al.*, *Genome.* 2002 Dec;45(6):1203-15).
- McCallum *et al.* (2000) *Nature Biotechnology* 18:455-457).



**This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**